

Maeotias inexspectata

Black Sea jellyfish

Threat Scores

- 1. Ecological Impact
 - The medusae eat larvae of an introduced mud crab, Rhithropanopeus harrisi (Mills and Sommer 1995)
 - Dense populations of this jelly can alter estuarine food webs
 - The prey of this jelly includes small adult crustaceans and larvae, so local populations of crabs and their relatives could be diminished
- 2. Invasive Potential
 - Know to be transported and discharged from ballast water
 - Tolerant of brackish and low salinity environments
 - Spread from original ports of entry into estuaries and river systems
- 3. Geographic Extent
 - Locally patchy
- 4. Management Difficulty
 - No effective eradication methods

Geography and Habitat

- 1. Native: Black Sea
- 2. Introduced: Atlantic Coast of United States, California
- 3. Habitats
 - Marine, brackish water, estuaries/bays

Invasion Pathways

1. Ballast water and sediments

Non-Native Locations

- 1. 41- Virginian
- 2. 42- Carolinian
- 3. 58- N. California

Sources

- 1. Molnar, Jennifer et al. 2008. Assessing the global threat of invasive species to marine biodiversity. Frontiers in ecology and the environment. Vol. 6, No. 9, pp. 485-492.
- 2. http://conserveonline.org/workspaces/global.invasive.assessment
- 3. http://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1066
- 4. http://www.elkhornslough.org/research/aquaticinvaders/aqua images/BLACKS~1.JPG

